

# Open Research Online

---

The Open University's repository of research publications  
and other research outputs

## Elevated InsP3R expression underlies enhanced calcium fluxes and spontaneous extra-systolic calcium release events in hypertrophic cardiac myocytes

### Journal Item

#### How to cite:

Harzheim, Dagmar; Talasila, Amarnath; Movassagh, Mehregan; Foo, Roger S-Y.; Figg, Nichola; Bootman, Martin D. and Roderick, H. Llewelyn (2010). Elevated InsP3R expression underlies enhanced calcium fluxes and spontaneous extra-systolic calcium release events in hypertrophic cardiac myocytes. *Channels (Austin)*, 4(1) pp. 67–71.

For guidance on citations see [FAQs](#).

© 2010 Landes Bioscience

Version: Version of Record

Link(s) to article on publisher's website:  
<http://dx.doi.org/doi:10.4161/chan.4.1.10531>

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

---

[oro.open.ac.uk](http://oro.open.ac.uk)

## Elevated InsP<sub>3</sub>R expression underlies enhanced calcium fluxes and spontaneous extra-systolic calcium release events in hypertrophic cardiac myocytes

Dagmar Harzheim,<sup>1,†</sup> Amarnath Talasila,<sup>1</sup> Mehregan Movassagh,<sup>2</sup> Roger S.-Y. Foo,<sup>2</sup> Nichola Figg,<sup>2</sup> Martin D. Bootman<sup>1</sup> and H. Llewelyn Roderick<sup>1,3,\*</sup>

<sup>1</sup>Department of Molecular Signaling; Babraham Institute; Cambridge, UK; <sup>2</sup>Addenbrooke's Centre for Clinical Investigation; Cambridge, UK;

<sup>3</sup>Department of Pharmacology; University of Cambridge; Cambridge, UK

<sup>†</sup>Current Address: Caesar Research Center; Bonn, Germany

Cardiac hypertrophy is associated with profound remodeling of Ca<sup>2+</sup> signaling pathways. During the early, compensated stages of hypertrophy, Ca<sup>2+</sup> fluxes may be enhanced to facilitate greater contraction, whereas as the hypertrophic heart decompensates, Ca<sup>2+</sup> homeostatic mechanisms are dysregulated leading to decreased contractility, arrhythmia and death. Although ryanodine receptor Ca<sup>2+</sup> release channels (RyR) on the sarcoplasmic reticulum (SR) intracellular Ca<sup>2+</sup> store are primarily responsible for the Ca<sup>2+</sup> flux that induces myocyte contraction, a role for Ca<sup>2+</sup> release via the inositol 1,4,5-trisphosphate receptor (InsP<sub>3</sub>R) in cardiac physiology has also emerged. Specifically, InsP<sub>3</sub>-induced Ca<sup>2+</sup> signals generated following myocyte stimulation with an InsP<sub>3</sub>-generating agonist (e.g., endothelin, ET-1), lead to modulation of Ca<sup>2+</sup> signals associated with excitation-contraction coupling (ECC) and the induction of spontaneous Ca<sup>2+</sup> release events that cause cellular arrhythmia. Using myocytes from spontaneously hypertensive rats (SHR), we recently reported that expression of the type 2 InsP<sub>3</sub>R (InsP<sub>3</sub>R2) is significantly increased during hypertrophy. Notably, this increased expression was restricted to the junctional SR in close proximity to RyRs. There, enhanced Ca<sup>2+</sup> release via InsP<sub>3</sub>Rs serves to sensitize neighboring RyRs causing an augmentation of Ca<sup>2+</sup> fluxes during ECC as well as an increase in non-triggered Ca<sup>2+</sup> release events.

Although the sensitization of RyRs may be a beneficial consequence of elevated InsP<sub>3</sub>R expression during hypertrophy, the spontaneous Ca<sup>2+</sup> release events are potentially of pathological significance giving rise to cardiac arrhythmia. InsP<sub>3</sub>R2 expression was also increased in hypertrophic hearts from patients with ischemic dilated cardiomyopathy and aortically-banded mice demonstrating that increased InsP<sub>3</sub>R expression may be a general phenomenon that underlies Ca<sup>2+</sup> changes during hypertrophy.

To meet increased haemodynamic requirements, or as a result of damage, the heart elicits a hypertrophic response. This response is characterized by an enlargement of the cardiac muscle mass due to increased cell size rather than cell number.<sup>1,2</sup> Hypertrophy results from both physiological (pregnancy, exercise) and pathological stimuli (hypertension, ischemic heart disease) and can initially be a beneficial adaptive response. However, persistent cardiac stress may lead to a decompensated form of hypertrophy, which can lead to arrhythmia and ultimately heart failure and death. Indeed, cardiac hypertrophy is a predictor of future heart failure and death.<sup>2,3</sup> These pathologies are of major clinical significance and together with other cardiovascular diseases are now the primary cause of death in the developed world.<sup>4</sup>

Ca<sup>2+</sup> is key to cardiac function—stimulating myocyte contraction via

**Key words:** IP3, IP3R, calcium, hypertrophy, cardiac, arrhythmia, hypertension.

Submitted: 10/17/09

Accepted: 10/17/09

Previously published online:  
[www.landesbioscience.com/journals/channels/article/10531](http://www.landesbioscience.com/journals/channels/article/10531)

\*Correspondence to: H. Llewelyn Roderick;  
Email: [Llewelyn.roderick@bbrsc.ac.uk](mailto:Llewelyn.roderick@bbrsc.ac.uk)

Addendum to: Harzheim D, Movassagh M, Foo RS, Ritter O, Tashfeen A, Conway SJ, Bootman MD, Roderick HL. Increased InsP<sub>3</sub>Rs in the junctional sarcoplasmic reticulum augment Ca<sup>2+</sup> transients and arrhythmias associated with cardiac hypertrophy. *Proc Natl Acad Sci USA* 2009; 106:11406–11; PMID: 19549843; DOI: 10.1073/pnas.0905485.

excitation-contraction coupling (ECC)<sup>5</sup> and controlling the transcription of genes that underlies the induction of hypertrophy.<sup>6,7</sup> ECC is initiated by an action potential (AP) originating at the sinoatrial node. Coordinated conduction of this action potential through the heart results in a synchronized contraction of the heart chambers. The propagating action potential depolarizes the myocyte plasma membrane thereby opening voltage-sensitive L-type  $\text{Ca}^{2+}$  channels and allowing  $\text{Ca}^{2+}$  to enter the cell. This 'trigger'  $\text{Ca}^{2+}$ , termed a sparklet,<sup>8</sup> induces opening of ryanodine receptor (RyR)  $\text{Ca}^{2+}$  release channels on the juxtaposed sarcoplasmic reticulum (SR) intracellular  $\text{Ca}^{2+}$  store—a process called  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release (CICR).<sup>9</sup> CICR causes a global elevation in  $\text{Ca}^{2+}$ , which leads to myocyte contraction. Relaxation is then brought about by extrusion of  $\text{Ca}^{2+}$  across the sarcolemma by the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger (NCX) and by its sequestration into the SR by the SERCA2  $\text{Ca}^{2+}$  ATPase.

In addition to RyRs, cardiomyocytes also express inositol 1,4,5-triphosphate receptor  $\text{Ca}^{2+}$  release channels ( $\text{InsP}_3\text{Rs}$ ), albeit outnumbered by RyRs 50:1.<sup>10–12</sup> Since  $\text{InsP}_3\text{Rs}$  have been discovered in the heart, it has been questioned how the relatively small amount of  $\text{Ca}^{2+}$  that is released via  $\text{InsP}_3$  can have any physiological function in the midst of the large amount of  $\text{Ca}^{2+}$  that is cycled through RyRs during every heart beat. In mammals, there are three  $\text{InsP}_3\text{R}$  isoforms (types 1, 2 and 3),<sup>13,14</sup> with type 2 being the main isoform in the heart.<sup>10,12</sup>  $\text{Ca}^{2+}$  release through  $\text{InsP}_3\text{Rs}$  has been shown to mediate inotropy and arrhythmogenic  $\text{Ca}^{2+}$  release events as well as contribute to the hypertrophic effect of  $\text{G}_q$ -coupled agonists such as the vasoactive peptide endothelin-1 (ET-1).<sup>15–22</sup>

Profound remodeling of the  $\text{Ca}^{2+}$  signaling apparatus has been reported to occur as a result of cardiac hypertrophy and failure.<sup>23</sup> Whilst enhancement of  $\text{Ca}^{2+}$  transients facilitates greater myocyte contraction during adaptive hypertrophy,  $\text{Ca}^{2+}$  fluxes are diminished during heart failure resulting in decreased contractility and thus, cardiac output.<sup>24</sup> Remodeling of the activity and function of the  $\text{Ca}^{2+}$  signaling proteome also underlies the increased

arrhythmias associated with hypertrophy and heart failure.<sup>25</sup> Whether altered  $\text{InsP}_3\text{R}$  signaling also contributes to remodeling of  $\text{Ca}^{2+}$  homeostasis during cardiac hypertrophy is not yet determined.

In a recent study,<sup>15</sup> we established that augmented expression and  $\text{Ca}^{2+}$  release through  $\text{InsP}_3\text{Rs}$  contribute to enhanced  $\text{Ca}^{2+}$  fluxes and increased spontaneous arrhythmogenic  $\text{Ca}^{2+}$  events associated with hypertrophy. As a model to study hypertrophy-dependent changes in  $\text{Ca}^{2+}$  signaling, we used hypertrophic ventricular cardiomyocytes from spontaneously hypertensive rats (SHR) and non-hypertrophic ventricular cardiomyocytes from Wistar Kyoto rats (WKY) as controls.<sup>26</sup> We found that in hypertrophic myocytes  $\text{InsP}_3\text{R2}$  expression was specifically increased in the junctional SR membrane, thereby increasing the numbers of  $\text{InsP}_3\text{Rs}$  in close proximity with type 2 RyRs (RyR2), the expression of which was unchanged as a result of hypertrophy. Enhanced  $\text{Ca}^{2+}$  release through these  $\text{InsP}_3\text{Rs}$  sensitized their neighboring RyRs, thereby enhancing CICR leading to bigger  $\text{Ca}^{2+}$  transients and greater myocyte contraction. Moreover, the sensitization of RyR opening through increased  $\text{InsP}_3$ -induced  $\text{Ca}^{2+}$  release (IICR) resulted in a higher propensity for the generation of elementary  $\text{Ca}^{2+}$  release events and extrasystolic  $\text{Ca}^{2+}$  transients, which could act as a substrate for cardiac arrhythmia *in vivo*. Interestingly, we could not only show that  $\text{InsP}_3\text{R2}$  expression was increased in hypertrophic rat cardiomyocytes but also in cardiomyocytes isolated from aortically-banded mice and in human hearts displaying ischemic dilated cardiomyopathy. We showed by immunoblotting and, as presented here, by immunohistochemistry that  $\text{InsP}_3\text{R2}$  expression is increased in diseased human hearts compared to control hearts (Fig. 1). These micrographs also illustrate that  $\text{InsP}_3\text{R2}$  expression is restricted to myocytes and is not present in non-myocyte cells, for example surrounding the vessels.


An increase in  $\text{InsP}_3\text{R2}$  expression therefore seems to be a general phenomenon occurring during cardiac disease. This is surprising considering the lack of consensus regarding the different effects of hypertrophy on  $\text{Ca}^{2+}$  signaling.<sup>23</sup>

Although amplification of  $\text{Ca}^{2+}$  cycling is considered as a general mechanism for increasing cardiac output during hypertrophy, myocytes isolated from certain hypertrophy models exhibit either no change, or even suppression, in their ECC-mediated  $\text{Ca}^{2+}$  fluxes.<sup>27,28</sup> As such, the mechanisms underlying remodeling of  $\text{Ca}^{2+}$  cycling during hypertrophy are not fully understood. In hypertrophic failing myocytes,  $\text{Ca}^{2+}$  cycling is suppressed causing decreased contractility. In this condition, defects in coupling between L-type channels and RyRs due to a breakdown in the t-tubular network,<sup>27</sup> decreases in SERCA-mediated  $\text{Ca}^{2+}$  sequestration into the SR lumen,<sup>29</sup> and increased  $\text{Ca}^{2+}$  leak from the SR via hyper-phosphorylated RyRs<sup>30</sup> have all been shown to be involved in pathology.

The reason for this array of  $\text{Ca}^{2+}$  signaling phenotypes during hypertrophy may lie in the panoply of hypertrophy models employed. For example, hypertrophy is induced by haemodynamic loading (such as aortic banding), interruption of myocardial blood supply (infarction) and interfering of heart rate (pacing or AV block).<sup>23,31</sup> More recently, genetic models of hypertrophy have also been employed.<sup>32</sup> Even more variability between studies is introduced by the hypertrophic stage at which  $\text{Ca}^{2+}$  signaling measurements are made and the location of the myocardium from which myocytes are isolated.<sup>23</sup> The SHR model offers considerable advantages. The temporal development of hypertrophy in SHRs has been well characterized and develops slowly between the age of 2 and 6 months subsequent to the onset of systemic hypertension.<sup>33</sup> After the development of hypertrophy, there is also a long latency before the development of heart failure at 20–24 months.<sup>26</sup> This contrasts with surgically-induced models where hypertrophy develops 1–2 weeks post surgery.<sup>31</sup> However, our results show that the increase in  $\text{InsP}_3\text{R}$  expression seems to be independent of how hypertrophy has been induced.<sup>15</sup>

Considering the pro-arrhythmic effect of increased  $\text{InsP}_3\text{R}$  expression, which is clearly evident in our study, this raises the question of why  $\text{InsP}_3\text{R}$  expression is enhanced in hypertrophic and failing cardiomyocytes? During hypertrophy a decrease in the coupling efficiency





**Figure 1.** Expression of InsP<sub>3</sub>R2 is increased in cardiomyocytes from human patients suffering from ischemic dilated cardiomyopathy. Immunohistochemical detection of InsP<sub>3</sub>R2 (brown) in cardiomyocytes from control patients (A) and in cardiomyocytes from patients showing ischemic dilated cardiomyopathy (B). Higher magnification images for (A and B) are presented in (C and D) respectively. Control staining of hypertrophic myocytes without using InsP<sub>3</sub>R2 primary antibody are presented at low and high magnification in (E and F) respectively. These data are representative of images taken from slides prepared from two control and two diseased hearts. Non-myocyte cells surrounding a blood vessel (indicated by the black arrow) do not stain for InsP<sub>3</sub>R2. The size of the bar is indicated in each image. The nuclei are indicated in blue.


between plasma membrane L-type channels and RyR channels due to an increase in the width of the dyadic cleft has been observed.<sup>28</sup> This deficit in ECC did not significantly affect contractile function and was overcome by beta-adrenergic stimulation. Here, sensitization of RyR opening due to increased Ca<sup>2+</sup> release through InsP<sub>3</sub>Rs could compensate for the lack of coupling efficiency, even under basal conditions.

What remains to be determined is how InsP<sub>3</sub>R2 expression during hypertrophy is regulated and how the striking junctional pattern of InsP<sub>3</sub>R2 expression

is achieved. Our data would suggest that InsP<sub>3</sub>R expression increases concomitant with or is a product of hypertrophy and is not causal. In support of this conclusion, non-hypertrophic hearts from four month old SHR do not exhibit increased InsP<sub>3</sub>R expression compared to WKY controls.<sup>15</sup> Moreover, hypertrophy is induced in vitro in an InsP<sub>3</sub>-dependent manner within 24 hours, a time period that would not allow for elevated InsP<sub>3</sub>R expression to have effect. Thus, it is likely that the role of increased InsP<sub>3</sub>Rs detected during hypertrophy is to modify calcium signals, and possibly to also act to sustain the

hypertrophic phenotype. The subcellular localization of InsP<sub>3</sub>R2 expression could be mediated via interactions with accessory proteins such as Ankyrin B, which has been shown to localize InsP<sub>3</sub>Rs.<sup>34</sup> Whether the mechanisms that mediate a preferential increase in InsP<sub>3</sub>Rs in the junctional zones and not around the nucleus during hypertrophy play any role in restricting the recently described pro-hypertrophic action of InsP<sub>3</sub>-induced calcium release (IICR) in the nucleus also remains to be determined.<sup>16,35</sup>

A dramatic consequence of hypertrophy for Ca<sup>2+</sup> signaling observed in our



**Figure 2.** Expression of  $ET_A R$  is increased in hypertrophic SHR hypertrophic myocytes. 25  $\mu$ g of total cell lysate prepared from left ventricle taken from SHR and WKY hearts from 6 month old animals was subjected to immunoblot analysis following SDS-PAGE. Analysis of samples prepared from 4 control (WKY) and 4 hypertrophic (SHR) hearts are shown. An immunoblot of  $\alpha$ -actinin in the same samples is presented as loading control. We have previously established that myocytes from 6 month old SHRs are hypertrophic exhibiting increased size and expression of atrial natriuretic factor (ANF).<sup>15</sup>

study was the increased occurrence of extra-systolic  $Ca^{2+}$  transients following ET-1 or  $InsP_3$  ester (cell permeant  $InsP_3$ ) application. The relatively low incidence of  $InsP_3$ -stimulated  $Ca^{2+}$  release events in WKY myocytes is consistent with our previous work in Wistar rat ventricular myocytes.<sup>10,20-22</sup> The enhanced incidence of extra-systolic  $Ca^{2+}$  transients in hypertrophic ventricular myocytes is reminiscent of previous findings in atrial myocytes, which express ~6-fold more  $InsP_3Rs$  under normal conditions.<sup>17,20,22,36</sup> Due to their higher basal expression of  $InsP_3Rs$ , stimulation of atrial myocytes with  $InsP_3$  or  $InsP_3$ -generating agonists, such as ET-1, potentially induce extra-systolic  $Ca^{2+}$  release events.<sup>20</sup> Although insignificant by themselves, these miniature  $Ca^{2+}$  release events can provide sufficient  $Ca^{2+}$  to catalyze further  $Ca^{2+}$  release through CICR and cause AP-independent increases in  $[Ca^{2+}]_i$  that could potentially result in arrhythmia.<sup>17</sup> This role of  $Ca^{2+}$  release through  $InsP_3Rs$  has been substantiated in  $InsP_3R2$ -deficient mice, in which ET-1- and  $InsP_3$ -stimulated arrhythmias in atrial myocytes were completely abolished.<sup>17</sup> Interestingly,  $InsP_3Rs$  in atrial myocytes are located on the SR underlying the sarcolemma<sup>20</sup>—the functional equivalent of the junctional zone created by the juxtaposition of the SR terminal cisternae and the t-tubular membrane in ventricular myocytes.<sup>37</sup> Thus, in hypertrophic ventricular myocytes, ECC may mirror that observed in atrial myocytes.

As well as being manifest as spontaneous  $Ca^{2+}$  release events and extra systolic  $Ca^{2+}$  transients, increased  $InsP_3R$  expression resulted in a greater rise in diastolic  $[Ca^{2+}]$  in hypertrophic myocytes following ET-1 or IICR stimulation.<sup>15</sup> As a result, diastolic  $[Ca^{2+}]$  was brought closer to the threshold for activation of RyRs, thereby increasing the incidence of stochastic  $Ca^{2+}$  release events and accelerating the generation of pacing-evoked  $Ca^{2+}$  transients. These findings are consistent with data from  $InsP_3R2$ -deficient atrial myocytes which exhibited a decrease in the ET-1-evoked elevation in diastolic  $[Ca^{2+}]$ .<sup>17</sup>

The arrhythmogenic effect of  $InsP_3$ -stimulated  $Ca^{2+}$  release events in the ventricles is potentially more catastrophic than in the atria. Whereas atrial fibrillation only has potentially serious consequences over the long term, ventricular fibrillation is potentially immediately life-threatening. This situation is further exacerbated by increases in the expression of additional components of the pathway that leads to  $InsP_3$  generation. Specifically, levels of cardiac ET-1 and also that of its receptor ( $ET_A R$ ), are elevated during development of hypertension and hypertrophy (Fig. 2).<sup>38,39</sup> Moreover, phospholipase C beta1 levels have also been recently reported to be increased during atrial hypertrophy.<sup>40</sup> Together, enhanced activity of this signaling pathway as a result of hypertrophy would promote greater  $Ca^{2+}$  release through  $InsP_3Rs$ , and heighten the

susceptibility for arrhythmias, even under non-stressed conditions.

Our recent study<sup>15</sup> presents a plausible mechanism why hypertrophic hearts are more prone to cardiac arrhythmia and proposes  $InsP_3R2$  as a possible target for the development of new drugs against hypertrophy-related arrhythmia.

## Materials and Methods

**Animal models.** Male SHRs and normotensive Wistar-Kyoto (WKY) rats were obtained from Harlan, UK, and were housed under control conditions with ad libitum food and water. All experiments were performed in accordance with the guidelines from the code of practice for humane killing under Schedule 1 of the Animals (Scientific Procedures) Act 1986.

**Patients.** Left ventricular tissue samples of human failing hearts were from individuals undergoing heart transplantation due to end-stage heart failure. All samples were obtained from male caucasians, aged 41–62. Samples from non-failing donor hearts were provided by the “UK Human Tissue Bank”. After cardiectomy, left ventricular samples were frozen in liquid nitrogen and stored at  $-80^{\circ}C$ . Detailed information about the patients is included in Harzheim et al. 2009. All experiments involving human tissue samples have been approved by the Cambridgeshire Research Ethics Committee.

**Immunohistochemistry.** Immunohistochemistry was performed as previously described.<sup>41</sup> Paraffin tissue sections were prepared from non-diseased human hearts and from hearts with ischemic dilated cardiomyopathy (tissue source described above). Sections were deparaffinized and antigen retrieved using basic citrate buffer, after which slides were washed with PBS and blocked with 0.3%  $H_2O_2$ /PBS for 10 mins. After a further wash in PBS, slides were incubated with 5% BSA/PBS for 1 hour at room temp. Slides were then incubated overnight at  $4^{\circ}C$  with polyclonal rat anti  $InsP_3R2$  antibody<sup>15</sup> diluted at 1:50 in 5% BSA. After removal of excess primary antibody by washing in PBS, slides were incubated with HRP-conjugated secondary antibody diluted at 1:400 for 30 minutes at room temp. After washing, slides were incubated with ABC reagent for 30 mins at



room temp after which they were washed, visualized with DAB, counterstained with Harris' haematoxylin and mounted.

**Immunoblot detection of ET<sub>A</sub>R.** Protein expression was detected as previously described.<sup>42</sup> In brief, proteins were extracted from ventricular myocytes from WKY rats and SHRs in lysis buffer (10 mM Tris-HCl, pH 7.6, 140 mM NaCl, 1 mM EDTA, 1% Triton X-100, protease and phosphatase inhibitor cocktail (Sigma)). Protein extracts (25 µg) were then subjected to SDS-PAGE (10% Bis-Tris NuPage, Invitrogen) and subsequent immunoblot analysis with antibodies directed against the ET<sub>A</sub>R (Alomone labs) and alpha-actinin (Sigma). Immune-complexes were detected by enhanced chemiluminescence with anti-mouse or anti-rabbit IgG coupled to horseradish peroxidase as the secondary antibody (Jackson Immuno Research Laboratories).

#### Acknowledgements

The work was supported by The BHF (PG/06/034/20637 and PG/07/040) and the BBSRC. H.L.R. is a Royal Society Research Fellow. D.H. also thanks Wolfson College for a JR Fellowship.

#### References

- Dorn GW, 2nd, Molkentin JD. Manipulating cardiac contractility in heart failure: data from mice and men. *Circulation* 2004; 109:20-2.
- Lorell BH, Carabello BA. Left ventricular hypertrophy: pathogenesis, detection and prognosis. *Circulation* 2000; 102:470-9.
- Vakili BA, Okin PM, Devereux RB. Prognostic implications of left ventricular hypertrophy. *American Heart Journal* 2001; 141:334-41.
- Allender S, Peto V, Scarborough P, Boxer A, Raynor M. Coronary Heart Disease Statistics. BHF, London 2006.
- Bers DM. Cardiac excitation-contraction coupling. *Nature* 2002; 415:198-205.
- Frey N, McKinsey TA, Olson EN. Decoding calcium signals involved in cardiac growth and function. *Nat Med* 2000; 6:1221-7.
- Heineke J, Molkentin JD. Regulation of cardiac hypertrophy by intracellular signaling pathways. *Nat Rev Mol Cell Biol* 2006; 7:589-600.
- Wang SQ, Song LS, Lakatta EG, Cheng H. Ca<sup>2+</sup> signaling between single L-type Ca<sup>2+</sup> channels and ryanodine receptors in heart cells. *Nature* 2001; 410:592-6.
- Roderick HL, Berridge MJ, Bootman MD. Calcium-induced calcium release. *Curr Biol* 2003; 13:425.
- Lipp P, Laine M, Tovey SC, Burrell KM, Berridge MJ, Li W, Bootman MD. Functional InsP<sub>3</sub> receptors that may modulate excitation-contraction coupling in the heart. *Curr Biol* 2000; 10:939-42.
- Moschella MC, Marks AR. Inositol 1,4,5-trisphosphate receptor expression in cardiac myocytes. *J Cell Biol* 1993; 120:1137-46.
- Perez PJ, Ramos-Franco J, Fill M, Mignery GA. Identification and functional reconstitution of the type 2 inositol 1,4,5-trisphosphate receptor from ventricular cardiac myocytes. *J Biol Chem* 1997; 272:23961-9.
- Foskett JK, White C, Cheung KH, Mak DO. Inositol trisphosphate receptor Ca<sup>2+</sup> release channels. *Physiol Rev* 2007; 87:593-658.
- Patel S, Joseph SK, Thomas AP. Molecular properties of inositol 1,4,5-trisphosphate receptors. *Cell Calcium* 1999; 25:247-64.
- Harzheim D, Movassagh M, Foo RS, Ritter O, Tashfeen A, Conway SJ, et al. Increased InsP<sub>3</sub>Rs in the junctional sarcoplasmic reticulum augment Ca<sup>2+</sup> transients and arrhythmias associated with cardiac hypertrophy. *Proc Natl Acad Sci USA* 2009; 106:11406-11.
- Higazi DR, Fearnley CJ, Drawnel FM, Talasila A, Corps EM, Ritter O, et al. Endothelin-1-stimulated InsP<sub>3</sub>-induced Ca<sup>2+</sup> release is a nexus for hypertrophic signaling in cardiac myocytes. *Mol Cell* 2009; 33:472-82.
- Li X, Zima AV, Sheikh F, Blatter LA, Chen J. Endothelin-1-induced arrhythmogenic Ca<sup>2+</sup> signaling is abolished in atrial myocytes of inositol-1,4,5-trisphosphate (IP<sub>3</sub>)-receptor type 2-deficient mice. *Circ Res* 2005; 96:1274-81.
- Luo D, Yang D, Lan X, Li K, Li X, Chen J, et al. Nuclear Ca<sup>2+</sup> sparks and waves mediated by inositol 1,4,5-trisphosphate receptors in neonatal rat cardiomyocytes. *Cell Calcium* 2008; 43:165-74.
- Luo DL, Gao J, Lan XM, Wang G, Wei S, Xiao RP, Han QD. Role of inositol 1,4,5-trisphosphate receptors in alpha1-adrenergic receptor-induced cardiomyocyte hypertrophy. *Acta Pharmacol Sin* 2006; 27:895-900.
- Mackenzie L, Bootman MD, Laine M, Berridge MJ, Thuring J, Holmes A, et al. The role of inositol 1,4,5-trisphosphate receptors in Ca<sup>2+</sup> signaling and the generation of arrhythmias in rat atrial myocytes. *J Physiol* 2002; 541:395-409.
- Proven A, Roderick HL, Conway SJ, Berridge MJ, Horton JK, Capper SJ, Bootman MD. Inositol 1,4,5-trisphosphate supports the arrhythmogenic action of endothelin-1 on ventricular cardiac myocytes. *J Cell Sci* 2006; 119:3363-75.
- Zima AV, Blatter LA. Inositol-1,4,5-trisphosphate-dependent Ca<sup>2+</sup> signaling in cat atrial excitation-contraction coupling and arrhythmias. *J Physiol* 2004; 555:607-15.
- Balke CW, Shorofsky SR. Alterations in calcium handling in cardiac hypertrophy and heart failure. *Cardiovasc Res* 1998; 37:290-9.
- Carvalho BM, Bassani RA, Franchini KG, Bassani JW. Enhanced calcium mobilization in rat ventricular myocytes during the onset of pressure overload-induced hypertrophy. *Am J Physiol Heart Circ Physiol* 2006.
- Kahan T, Bergfeldt L. Left ventricular hypertrophy in hypertension: its arrhythmogenic potential. *Heart* 2005; 91:250-6.
- Bing OH, Brooks WW, Robinson KG, Slawsky MT, Hayes JA, Litwin SE, et al. The spontaneously hypertensive rat as a model of the transition from compensated left ventricular hypertrophy to failure. *J Mol Cell Cardiol* 1995; 27:383-96.
- Gomez AM, Valdivia HH, Cheng H, Lederer MR, Santana LF, Cannell MB, et al. Defective excitation-contraction coupling in experimental cardiac hypertrophy and heart failure. *Science* 1997; 276:800-6.
- Xu M, Zhou P, Xu SM, Liu Y, Feng X, Bai SH, et al. Intermolecular failure of L-type Ca<sup>2+</sup> channel and ryanodine receptor signaling in hypertrophy. *PLoS Biol* 2007; 5:21.
- Kubalova Z, Terentyev D, Viatchenko-Karpinski S, Nishijima Y, Gyorke I, Terentyeva R, et al. Abnormal intrastore calcium signaling in chronic heart failure. *Proc Natl Acad Sci USA* 2005; 102:14104-9.
- Wehrens XHT, Lehnart SE, Reiken S, Vest JA, Wronska A, Marks AR. Ryanodine receptor/calcium release channel PKA phosphorylation: A critical mediator of heart failure progression. *Proc Natl Acad Sci USA* 2006; 103:511-8.
- Balakumar P, Singh AP, Singh M. Rodent models of heart failure. *J Pharmacol Toxicol Meth* 2007; 56:1-10.
- Wang Y. Signal transduction in cardiac hypertrophy—dissecting compensatory versus pathological pathways utilizing a transgenic approach. *Curr Opin Pharmacol* 2001; 1:134-40.
- Bell D, Kelso EJ, Argent CCH, Lee GR, Allen AR, McDermott BJ. Temporal characteristics of cardiomyocyte hypertrophy in the spontaneously hypertensive rat. *Cardiovasc Pathol* 2004; 13:71-8.
- Mohler PJ, Davis JQ, Bennett V. Ankyrin-B Coordinates the Na/K ATPase, Na/Ca Exchanger, and InsP<sub>3</sub> Receptor in a Cardiac T-Tubule/SR Microdomain. *PLoS Biology* 2005; 3:423.
- Wu X, Zhang T, Bossuyt J, Li X, McKinsey TA, Dedman JR, et al. Local InsP<sub>3</sub>-dependent perinuclear Ca<sup>2+</sup> signaling in cardiac myocyte excitation-contraction coupling. *J Clin Invest* 2006; 116:675-82.
- Kocksamper J, Seidlmayer L, Walther S, Hellenkamp K, Maier LS, Pieske B. Endothelin-1 enhances nuclear Ca<sup>2+</sup> transients in atrial myocytes through Ins(1,4,5)P<sub>3</sub>-dependent Ca<sup>2+</sup> release from perinuclear Ca<sup>2+</sup> stores. *J Cell Sci* 2008; 121:186-95.
- Bootman MD, Higazi DR, Coombes S, Roderick HL. Calcium signaling during excitation-contraction coupling in mammalian atrial myocytes. *J Cell Sci* 2006; 119:3915-25.
- Iyer RS, Singh G, Rebello S, Roy S, Bhat R, Vidyasagar D, Gulati A. Changes in the concentration of endothelin-1 during development of hypertensive rats. *Pharmacology* 1995; 51:96-104.
- Zolk O, Quatteck J, Seeland U, El-Armouche A, Eschenhagen T, Bohm M. Activation of the cardiac endothelin system in left ventricular hypertrophy before onset of heart failure in TG(mREN2)27 rats. *Cardiovasc Res* 2002; 53:363-71.
- Woodcock EA, Grubb DR, Filtz TM, Marasco S, Luo J, McLeod-Dryden TJ, et al. Selective activation of the "b" splice variant of phospholipase Cbeta1 in chronically dilated human and mouse atria. *J Mol Cell Cardiol* 2009.
- Clarke MC, Figg N, Maguire JJ, Davenport AP, Goddard M, Littlewood TD, Bennett MR. Apoptosis of vascular smooth muscle cells induces features of plaque vulnerability in atherosclerosis. *Nat Med* 2006; 12:1075-80.
- Kasri NN, Holmes AM, Bultynck G, Parys JB, Bootman MD, Rietdorf K, et al. Regulation of InsP<sub>3</sub> receptor activity by neuronal Ca<sup>2+</sup>-binding proteins. *EMBO J* 2004; 23:312-21.